



BRAFTOVI™ (encorafenib) + MEKTOVI® (binimetinib) Receives Positive CHMP Opinion for Advanced BRAF-mutant Melanoma

July 27, 2018

- **Positive CHMP opinion is based on Phase 3 COLUMBUS trial which demonstrated nearly 15 months median progression-free survival and over 30 months median overall survival -**
- **U.S. FDA recently approved BRAFTOVI + MEKTOVI for advanced BRAF-mutant melanoma -**

BOULDER, Colo., July 27, 2018 /PRNewswire/ -- Array BioPharma Inc. (NASDAQ: ARRY) today announced that the Committee for Medicinal Products for Human Use (CHMP) of the European Medicines Agency (EMA) adopted a positive opinion recommending approval of BRAFTOVI™ in combination with MEKTOVI® for the treatment of adult patients with unresectable or metastatic melanoma with a *BRAF*^{V600} mutation. The CHMP recommendation will now be reviewed by the European Commission (EC), which has the authority to approve medicines for the European Union (EU). The final EC decision, expected by the end of September, will be applicable to all 28 EU member states, as well as Liechtenstein, Iceland and Norway.

"Following the recent U.S. FDA approval of BRAFTOVI + MEKTOVI for advanced *BRAF*-mutant melanoma, we are pleased to move one step closer to European approval," said Ron Squarer, Chief Executive Officer. "We are proud that the combination of BRAFTOVI + MEKTOVI represents a new standard of care for *BRAF*-mutant melanoma patients in critical need of additional treatment options."

The positive CHMP opinion is based on results from the Phase 3 COLUMBUS trial, which demonstrated that the combination BRAFTOVI + MEKTOVI achieved a median progression-free survival (mPFS) of nearly 15 months [14.9 months versus vemurafenib monotherapy at 7.3 months; hazard ratio (HR) 0.54 (95% CI, 0.41–0.71), *p*<0.0001].

In June 2018, Array also [announced](#) updated results from the COLUMBUS trial, which demonstrated that BRAFTOVI + MEKTOVI was the first targeted therapy to achieve over 30 months overall survival (OS) in a Phase 3 trial and reduced the risk of death compared to treatment with vemurafenib [HR (0.61), (95% CI 0.47-0.79, *p* <0.0001)]. Median OS was 33.6 months for patients treated with the combination, compared to 16.9 months for patients treated with vemurafenib as a monotherapy.

The most common grade 3-4 adverse events seen in more than 5% of patients were increased gamma-glutamyltransferase (9%), increased blood creatine phosphokinase (7%) and hypertension (6%).

In the U.S., BRAFTOVI + MEKTOVI are approved for the treatment of unresectable or metastatic melanoma with a *BRAF*^{V600E} or *BRAF*^{V600K} mutation, as detected by an FDA-approved test. BRAFTOVI is not indicated for treatment of patients with wild-type *BRAF* melanoma.

Only 5% of patients who received BRAFTOVI + MEKTOVI discontinued treatment due to adverse reactions. The most common adverse reactions (≥25%) in patients receiving BRAFTOVI + MEKTOVI were fatigue, nausea, diarrhea, vomiting, abdominal pain, and arthralgia.

Detailed recommendations for the use of these products will be described in the summary of product characteristics (SmPC), which will be published in the European public assessment report (EPAR) and made available in all official EU languages after the marketing authorization has been granted by the EC. Pierre Fabre has exclusive rights to commercialize both products in Europe, Asia and Latin America.

About BRAF-mutant Metastatic Melanoma

Melanoma develops when unrepaired DNA damage to skin cells triggers mutations that may lead them to multiply and form malignant tumors. Metastatic melanoma is the most serious and life-threatening type of skin cancer and is associated with low survival rates. [1,2] There are a variety of gene mutations that can lead to metastatic melanoma. The most common genetic mutation in metastatic melanoma is *BRAF*. There are about 200,000 new cases of melanoma diagnosed worldwide each year, approximately half of which have *BRAF* mutations, a key target in the treatment of metastatic melanoma. [1-5]

About BRAFTOVI + MEKTOVI

BRAFTOVI is an oral small molecule BRAF kinase inhibitor and MEKTOVI is an oral small molecule MEK inhibitor which target key enzymes in the MAPK signaling pathway (RAS-RAF-MEK-ERK). Inappropriate activation of proteins in this pathway has been shown to occur in many cancers including melanoma, colorectal cancer, non-small cell lung cancer, thyroid and others. In the U.S., BRAFTOVI + MEKTOVI are approved for the treatment of unresectable or metastatic melanoma with a *BRAF*^{V600E} or *BRAF*^{V600K} mutation, as detected by an FDA-approved test. BRAFTOVI is not indicated for treatment of patients with wild-type *BRAF* melanoma.

Array has exclusive rights to BRAFTOVI and MEKTOVI in the U.S. and Canada. Array has granted Ono Pharmaceutical exclusive rights to commercialize both products in Japan and South Korea, Medison exclusive rights to commercialize both products in Israel and Pierre Fabre exclusive rights to commercialize both products in all other countries, including Europe, Asia and Latin America.

BRAFTOVI and MEKTOVI are not approved outside of the United States. The European Medicines Agency (EMA), the Swiss Medicines Agency (Swissmedic) and the Australian Therapeutic Goods Administration (TGA) are currently reviewing the Marketing Authorization Applications, and the Pharmaceuticals and Medical Devices Agency (PMDA) in Japan is currently reviewing the Manufacturing and Marketing Approval Applications for BRAFTOVI and MEKTOVI.

About COLUMBUS

The COLUMBUS trial (NCT01909453) is a two-part, international, randomized, open label Phase 3 trial evaluating the efficacy and safety of BRAFTOVI (encorafenib) in combination with MEKTOVI (binimetinib) compared to vemurafenib and encorafenib monotherapy in 921 patients with locally advanced, unresectable or metastatic melanoma with *BRAF*^{V600} mutation. All secondary efficacy analyses, including overall survival, are descriptive in nature. Over 200 sites across North America, Europe, South America, Africa, Asia and Australia participated in the trial.

Indications and Usage

BRAFTOVI™ (encorafenib) and MEKTOVI® (binimetinib) are kinase inhibitors indicated for use in combination for the treatment of patients with unresectable or metastatic melanoma with a *BRAF*^{V600E} or *BRAF*^{V600K} mutation, as detected by an FDA-approved test.

Limitations of Use: BRAFTOVI is not indicated for the treatment of patients with wild-type *BRAF* melanoma.

BRAFTOVI + MEKTOVI Important Safety Information

The information below applies to the safety of the combination of BRAFTOVI and MEKTOVI unless otherwise noted.

Warnings and Precautions

New Primary Malignancies: New primary malignancies, cutaneous and non-cutaneous malignancies can occur. In the COLUMBUS trial, cutaneous squamous cell carcinoma, including keratoacanthoma, occurred in 2.6% and basal cell carcinoma occurred in 1.6% of patients. Perform dermatologic evaluations prior to initiating treatment, every 2 months during treatment, and for up to 6 months following discontinuation of treatment. Discontinue BRAFTOVI for *RAS* mutation-positive non-cutaneous malignancies.

Tumor Promotion in *BRAF* Wild-Type Tumors: Confirm evidence of *BRAF*^{V600E} or *BRAF*^{V600K} mutation prior to initiating BRAFTOVI.

Cardiomyopathy: In the COLUMBUS trial, cardiomyopathy occurred in 7% and Grade 3 left ventricular dysfunction occurred in 1.6% of patients. Cardiomyopathy resolved in 87% of patients. Assess left ventricular ejection fraction by echocardiogram or MUGA scan prior to initiating treatment, 1 month after initiating treatment, and then every 2 to 3 months during treatment. The safety has not been established in patients with a baseline ejection fraction that is either below 50% or below the institutional lower limit of normal.

Venous Thromboembolism (VTE): In the COLUMBUS trial, VTE occurred in 6% of patients, including 3.1% of patients who developed pulmonary embolism.

Hemorrhage: In the COLUMBUS trial, hemorrhage occurred in 19% of patients and ≥Grade 3 hemorrhage occurred in 3.2% of patients. Fatal intracranial hemorrhage in the setting of new or progressive brain metastases occurred in 1.6% of patients.

Ocular Toxicities: In the COLUMBUS trial, serous retinopathy occurred in 20% of patients; 8% were retinal detachment and 6% were macular edema. Symptomatic serous retinopathy occurred in 8% of patients with no cases of blindness. In patients with *BRAF* mutation-positive melanoma across multiple clinical trials, 0.1% of patients experienced retinal vein occlusion (RVO). Permanently discontinue MEKTOVI in patients with documented RVO. In COLUMBUS, uveitis, including iritis and iridocyclitis, was reported in 4% of patients. Assess for visual symptoms at each visit. Perform ophthalmic evaluation at regular intervals and for any visual disturbances.

Interstitial Lung Disease (ILD): ILD, including pneumonitis, occurred in 0.3% of patients with *BRAF* mutation-positive melanoma across multiple clinical trials. Assess new or progressive unexplained pulmonary symptoms or findings for possible ILD.

Hepatotoxicity: In the COLUMBUS trial, the incidence of Grade 3 or 4 increases in liver function laboratory tests was 6% for alanine aminotransferase (ALT) and 2.6% for aspartate aminotransferase (AST). Monitor liver laboratory tests before and during treatment and as clinically indicated.

Rhabdomyolysis: In the COLUMBUS trial, elevation of laboratory values of serum creatine phosphokinase (CPK) occurred in 58% of patients. Rhabdomyolysis was reported in 0.1% of patients with *BRAF* mutation-positive melanoma across multiple clinical trials. Monitor CPK periodically and as clinically indicated.

QTc Prolongation: In the COLUMBUS trial, an increase in QTcF to >500 ms was measured in 0.5% (1/192) of patients. Monitor patients who already have or who are at significant risk of developing QTc prolongation. Correct hypokalemia and hypomagnesemia prior to and during BRAFTOVI administration. Withhold, reduce dose, or permanently discontinue for QTc >500 ms.

Embryo-Fetal Toxicity: BRAFTOVI or MEKTOVI can cause fetal harm when administered to pregnant women. Nonhormonal contraceptives should be used during treatment and for at least 30 days after the final dose for patients taking BRAFTOVI + MEKTOVI.

Adverse Reactions

The most common adverse reactions (≥20%, all Grades, in the COLUMBUS trial) were: fatigue, nausea, diarrhea, vomiting, abdominal pain, arthralgia, myopathy, hyperkeratosis, rash, headache, constipation, visual impairment, serous retinopathy.

In the COLUMBUS trial, the most common laboratory abnormalities (≥20%, all Grades) included: increased creatinine, increased CPK, increased gamma glutamyl transferase, anemia, increased ALT, hyperglycemia, increased AST, and increased alkaline phosphatase.

Drug interactions

Avoid concomitant use of strong or moderate CYP3A4 inhibitors or inducers and sensitive CYP3A4 substrates with BRAFTOVI. Modify BRAFTOVI dose if concomitant use of strong or moderate CYP3A4 inhibitors cannot be avoided.

Please see full Prescribing Information for BRAFTOVI and full Prescribing Information for MEKTOVI for additional information. You may report side effects to the FDA at (800) FDA-1088 or www.fda.gov/medwatch. You may also report side effects to Array at 1-844-Rx-Array (1-844-792-7729).

About Array BioPharma

Array BioPharma Inc. is a fully-integrated, biopharmaceutical company focused on the discovery, development and commercialization of safe and effective targeted small molecule drugs to treat patients afflicted with cancer and other conditions. Array markets in the United States BRAFTOVI™ capsules in combination with MEKTOVI® tablets for the treatment of patients with unresectable or metastatic melanoma with a *BRAF*^{V600E} or *BRAF*^{V600K} mutation. Array's lead clinical programs, encorafenib and binimetinib, are being investigated in over 30 clinical trials across a number of solid tumor indications, including a Phase 3 trial in *BRAF*-mutant colorectal cancer. Array's pipeline includes several additional advanced programs including selumetinib (partnered with AstraZeneca), larotrectinib (partnered with Loxo Oncology), ipatasertib (partnered with Genentech), tucatinib (partnered with Seattle Genetics) and ARRY-797 (being developed by Yarra Therapeutics, a wholly-owned subsidiary of Array), all of which are

currently in registration trials. Ganovo® (danoprevir, partnered with Roche) was recently approved in China for the treatment of viral hepatitis C. For more information on Array, please visit www.arraybiopharma.com or follow @arraybiopharma on Twitter and LinkedIn.

References

- [1] Melanoma Skin Cancer. American Cancer Society. Available at: <https://www.cancer.org/cancer/melanoma-skin-cancer.html>. Accessed January 2018.
- [2] A Snapshot of Melanoma. National Cancer Institute. Available at: <https://seer.cancer.gov/statfacts/html/melan.html>. Accessed January 2018.
- [3] Globocan 2012: Estimated Cancer Incidence, Mortality and Prevalence Worldwide in 2012. http://globocan.iarc.fr/Pages/fact_sheets_population.aspx. Accessed January 2018.
- [4] Klein O, et al. *Eur J Cancer*, 2013.
- [5] American Cancer Society. What Causes Melanoma Skin Cancer? 2016. <https://www.cancer.org/cancer/melanoma-skin-cancer/causes-risks-prevention/what-causes.html>. Accessed April 11, 2018.
- [6] BRAFTOVI™ (encorafenib) Prescribing Information Array BioPharma Inc., June 2018
- [7] MEKTOVI® (binimetinib) Prescribing Information. Array BioPharma Inc., June 2018

Array BioPharma Forward-Looking Statement

This press release contains forward-looking statements within the meaning of the Private Securities Litigation Reform Act of 1995, including, among others, statements about the future development plans of encorafenib and binimetinib; expectations that events will occur that will create greater value for Array; and the potential for the results of current and future clinical trials to support regulatory approval or the marketing success of encorafenib and binimetinib. Because these statements reflect our current expectations concerning future events and involve significant risks and uncertainties, our actual results could differ materially from those anticipated in these forward-looking statements as a result of many factors. These factors include, but are not limited to, the potential that the FDA, EMA or other regulatory agencies determine results from clinical trials are not sufficient to support registration or marketing approval of encorafenib and binimetinib; our ability to effectively and timely conduct clinical trials in light of increasing costs and difficulties in locating appropriate trial sites and in enrolling patients who meet the criteria for certain clinical trials; risks associated with our dependence on third-party service providers to successfully conduct clinical trials and to manufacture drug substance and product within and outside the U.S.; our ability to grow and successfully develop commercialization capabilities; our ability to achieve and maintain profitability and maintain sufficient cash resources; and our ability to attract and retain experienced scientists and management. Additional information concerning these and other risk factors can be found in our most recent annual report filed on Form 10-K, in our quarterly reports filed on Form 10-Q, and in other reports filed by Array with the Securities and Exchange Commission. We are providing this information as of July 27, 2018. We undertake no duty to update any forward-looking statements to reflect the occurrence of events or circumstances after the date of such statements or of anticipated or unanticipated events that alter any assumptions underlying such statements.

BRAFTOVI™ is a trademark of Array BioPharma Inc.

MEKTOVI® is a registered trademark of Array BioPharma Inc. in the United States and various other countries.

CONTACTS:

Investor Relations

Array BioPharma

Andrea N. Flynn, Ph.D.

Senior Director, Investor Relations & Corporate Communications

(303) 381-6600

ir@arraybiopharma.com

Media

Y&R PR

Erika Hackmann, Media Relations

(917) 538-3375

erika.hackmann@yr.com



View original content with multimedia: <http://www.prnewswire.com/news-releases/braftovi-encorafenib--mektovi-binimetinib-receives-positive-chmp-opinion-for-advanced-braf-mutant-melanoma-300687771.html>

SOURCE Array BioPharma Inc.